

1003 E FLORIDA AVE. SUITE 101 **HEMET, CA 92543** PH: (951) 652-2252 FAX:(951)658-6476

MILAN S CHAKRABARTY, M.D. SANDRA DEL VALLE, PA-C

Tima.

Date:

TAHIR QASEEM, MD. F.A.C.P., F.A.C.G. YAHAIRA SANCHEZ, FNP-C

PATIENT INSTRUCTI	ONS FOR UNDERGOING SMALL BOWEL CAPSULE ENDOSCOPY
Patient Information	
First Name:	Last Name:
Date of Birth:	Age:
SMALL BOWEL CAPSULE ENDO	SCOPY
capsule endoscopy. This procedure investions through your digestive system while to images. The Data Recorder is worn over processing. The NaviCam SB capsule is	a part of your medical evaluation you should undergo an examination known as small bowel (SB) olves ingesting a small (the size of a large vitamin pill) NaviCam SB capsule which will pass naturally aking pictures of the intestine. The images are transmitted to the DataRecorder which saves all the er your clothes on a Crossbody Belt. After 8 hours the Crossbody Belt will be removed for disposable and will be excreted naturally in your bowel movement. In the rare case that it will not be removed endoscopically or surgically. For your physician to get the most accurate information follow the instructions below:
The Day before Capsule Endos	сору

- 1. After lunch (12:00 p.m.) on the day before the SB capsule endoscopy examination, start a clear liquid diet. (See page 2 for the approved list of clear liquids)
- 2. At approximately 6:00 p.m. mix Miralax 119g into Gatorade 28 oz (no red, purple or orange). Drink 8 oz every 10-15 minutes until completed. Stop drinking fluids at 10:00 p.m.
- 3. Any necessary medications must be taken no later than 6:00 a.m. on the day of your procedure with a small amount of water.

The Day of Capsule Endoscopy

- 1. Arrive at the ofice at 8:00 a.m.
- 2. Wear loose fitting, two-piece clothing. Please wear a shirt made of thin material such as a cotton t-shirt.
- 3. At the office you will have a pre-procedure interview. During this interview you should be informed that the SB capsule endoscopy is associated with a small chance of intestinal obstruction.
- 4. The Crossbody Belt will be applied to your abdomen on top of your clothing. It will be connected to the DataRecorder. After that you will ingest the NaviCam SB capsule.

After Ingesting the NaviCam SB Capsule

- 1. After ingesting the NaviCam SB capsule, you may not have anything to eat or drink until 10:00 a.m. At 10:00 a.m. you may have water only.
- 2. Starting at 12:00 p.m. you may resume a clear liquid diet until you return to the office and have the equipment removed. You will return to the office at 4:30 p.m.
- 3. After ingesting the NaviCam capsule and until it is excreted, you should not be near any source of powerful electromagnetic field such as one created near an MRI device or amateur (ham) radio. Occasionally, some images may be lost due to radio interference. In rare occasions this may result in the need to repeat the SB capsule endoscopy examination.
- 4.SB capsule endoscopy lasts approximately 8 hours and is considered complete according to your physician's instructions. Do not disconnect the equipment or remove the belt at anytime during this period. Since the Data Recorder is actually a small computer it should be treated with utmost care and protection. Avoid sudden movement and banging of the Data Recorder.
- 5. During the SB capsule endoscopy, you will need to verify the light on the Data Recorder is green.

 If, for some reason stops, contact your physician's office.
- 6. Avoid strenuous physical activity such as exercising.

After completing the SB Capsule Endoscopy

1. If you did not positively verify the excretion of the NaviCam SB capsule from your body, and you develop unexplained post procedure nausea, abdominal pain or vomiting, contact your physician's office for evaluation and possible abdominal X-ray.

Clear Liquid Diet

- Coffee (black or with sugar), tea, soft drinks
- Strained fruit juices (apple, white grape), Lemonade (from a mix)
- Clear broth
- Gelatin (Jell-O), popsicles (no red, purple, or orange)
- Sugar, Honey, Syrup
- Clear Hard candy